

# Insights in Veterinary Science

Volume - 6, Issue - 1

**Letter to Editor**      **Published Date:-2022-09-14 00:00:00**

[The therapeutic impact of the interaction between horses and patients with mental disorders: Veterinary and psychodynamic reflexions in the clinical application of equine assisted therapy](#)

An increased scientific interest is observed nowadays in the clinical application of Equine Assisted therapy for patients with mental disorders, reflected on recent numerous systematic reviews, scoping reviews and concept analyses.

---

**Review Article**      **Published Date:-2022-04-29 11:21:27**

[Review on epidemiology of bovine hemoparasites in Ethiopia](#)

A literature-based review was made to assess available information on bovine hemoparasites related to their epidemiology, distribution, and economic importance in Ethiopia. Babesiosis, anaplasmosis, cowdriosis, theileriosis, and trypanosomosis are the major hemo-parasitic disease of bovine in Ethiopia. Their adverse effects on the health of the animals can decrease production and productivity. Hemoparasites generally cause fever, anemia, jaundice, anorexia, weight loss, milk drop, malaise, swelling of lymph nodes, dyspnoea, diarrhea, nervous disorders, and death by affecting blood vessels and/or lymphatic system of the animal. Reports from different parts of the country displayed there is a high distribution of bovine hemoparasitic disease throughout the country. Anaplasmosis, Babesiosis (redwater), Ehrlichiosis (Heartwater), Theileriosis, and Trypanosomosis are the major hemoparasitic diseases with heavy economic losses. Their mode of transmission was by arthropod vectors ticks and flies. Applying effective vector control and using vaccines drugs are the two most important control methods for hemoparasites diseases. Also having knowledge of parasite life cycles, their biological vector, and the immune response of bovines to vectors and parasites were also used in the successful application of control strategies. Creating awareness of the mode of transmission, method of control, and prevention of hemoparasitic disease of bovine to livestock owners were warranted to decrease the effect of the disease.

---

**Review Article**      **Published Date:-2022-03-15 10:35:09**

[Fish-borne parasites proficient in zoonotic diseases: a mini review](#)

Fish-borne parasitic zoonoses are primarily found in people living in developing and underdeveloped countries. The parasites that cause such zoonoses like Trichinella and Taenia are well-known in developed nations, but few people are familiar with fish-borne parasitic zoonoses, which are largely caused by helminths and protozoans. In general, parasitic zoonoses transmitted by fish are rarely life-threatening, although cases and reports of such infestations have increased over the world. The list of parasitic organisms is extensive. This article attempts to provide a complete overview of the many fish-borne parasites that can cause zoonosis among humans and animals alike.

---

**Research Article**      **Published Date:-2022-01-12 09:25:42**

[Collection and evaluation of indigenous buck semen at the coastal region of Bangladesh](#)

---

Evaluation of semen characteristics is an important and prior for semen preservation. The aim of this study was to collection and evaluation of indigenous buck semen in the coastal region of Bangladesh. The semen was collected from bucks through artificial vagina method. The colour, odour, volume, viscosity, mass activity, consistency, concentration and individual sperm motility were analysed and recorded after collection from pre-selected four bucks. The colour and odour of all buck (B) semen were creamy white to milky white and fishy smell, respectively. In this study, we found that the average volume of B-1, B-2, B-3 and B-4 were 0.74, 0.98, 0.42 and 0.60 ml, respectively. The average grading of viscosity of B-1, B-2, B-3 and B-4 were 3.2, 3.8, 2.6 and 3.0, respectively. The average grading of mass activity of B-1, B-2, B-3 and B-4 were 3.6, 3.2, 2.4 and 3.4, respectively. The consistency of B-1, B-2, B-3 and B-4 were 4.2, 4.8, 2.8 and 4.0, respectively. The concentration of B-1, B-2, B-3 and B-4 were 1.58, 1.94, 0.62 and 1.54  $\times 10^9$  per ml of semen volume. The average percentage of individual sperm motility of B-1, B-2, B-3 and B-4 were 81, 71, 66 and 80%, respectively. Viscosity, mass activity, consistency, concentration and individual sperm motility were significantly ( $p < 0.05$ ) correlate with each other. It may be concluded that the data about semen of the bucks are in acceptable level for preservation. Further study will be designed for the evaluation of viability and motility of sperm before and after freezing as liquid semen.

---